

A small-scale bacterial-based liquid Culture Method for *Steinernema hermaphroditum*

Nathan Y. Rodak^{1,2}, Chieh-Hsiang Tan^{1§}, Paul W. Sternberg^{1§}

¹Division of Biology and Biological Engineering, California Institute of Technology, 1200 East California Boulevard, Pasadena, CA, USA

²Current Address: Case Western Reserve University, Cleveland, OH, USA

[§]To whom correspondence should be addressed: chtan@caltech.edu; pws@caltech.edu

Abstract

Entomopathogenic nematodes (EPN) infect and kill their insect host with the help of symbiotic bacteria. The only known hermaphroditic (androdiecious) EPN, the clade IV *Steinernema hermaphroditum*, offers opportunities for exploring both parasitic and mutualistic symbiosis, as well as for evolutionary and developmental studies. Experimental and genetic analysis of this animal is now facilitated through the development of forward and reverse genetic tools and improved culturing techniques. Here, we describe a liquid-culture technique adapted for this worm. The method can be a starting point for the development of large-scale cultivation of the worm and provides a method to generate infective juveniles without an insect host and either with or without its native symbiotic bacteria.




Figure 1. A small-scale bacterial-based liquid Culture Method for *Steinernema hermaphroditum*:

(A) A gravid adult *S. hermaphroditum* grown at 25°C in S Medium liquid culture using *C. aquatica* as a food source. Left: anterior; down: ventral. Scale bar, 100 μm . (B, C) Bar charts showing the number of days required for P_0 embryos to develop into adults and produce F_1 larvae (upper charts, the day when the first F_1 larvae appear) and for the development of infective juveniles (IJs) (lower charts, the day when the first IJs appear). Three independent populations of worms were assayed for each condition which gives the same results. (B) Liquid culture in S Medium containing 20%, 50%, or 100% of concentrated *C. aquatica* suspension in S Medium by volume. (C) Liquid culture in S Medium containing 5%, 10%, or 20% (by volume) of concentrated *X. griffinae* suspension in S Medium. Worms failed to produce progeny when grown in 5% *X. griffinae*. (D, E) Worms grow faster in cultures with a higher concentration of food. Average lengths of worms grown in liquid culture containing varying concentrations of *C. aquatica* suspension. (D) 0% - 20% of concentrated *C. aquatica* by volume. Values are mean \pm SD ($n = 8 - 50$). (E). 0% - 50% of concentrated *C. aquatica* by volume. Data from three independent populations are presented separately. Values are mean \pm SD ($n = 5 - 15$, 15 worms were measured in most data points). (F) An infective juvenile (IJ) stage *S. hermaphroditum* from a S Medium / *C. aquatica* liquid culture incubated at 25°C. The worm has a sealed mouth, a constricted pharynx, and the entire body is enclosed by a seemingly detached second larval stage (J2) cuticle (red arrows) that is highly visible. Scale bar, 100 μm .

Description

A liquid-based culturing system for small animals such as *Caenorhabditis elegans* (Maupas 1900) is helpful in that it provides not only a method for the mass production of the animal but also is a valuable research tool, providing a uniform environment as a population grows. Liquid culture has been useful in the study of aspects of *C. elegans* biology, including stress resistance, induction of dauer larvae, and lifespan (Boyd et al. 2003; Baugh et al. 2011; Park et al. 2017; Hibshman et al. 2021). *Steinernema hermaphroditum* (Stock et al. 2004) is a Clade IV nematode that is evolutionarily distinct from the Clade V nematodes currently used extensively in the laboratory, such as *C. elegans* (Corsi et al. 2015), *Oscheius tipulae* (Lam and Webster 1971; Felix 2006), and *Pristionchus pacificus* (Sommer et al. 1996; Sommer 2006). It is also the only characterized consistently hermaphroditic entomopathogenic nematode (EPN) (Cao et al. 2022). These characteristics, along with a chromosomally scaffolded annotated genome and the availability of both forward and reverse genetic techniques, establish *S. hermaphroditum* as an excellent model for studying evolution, development, mutualistic symbiosis, and parasitism (Cao et al. 2022; Cao 2023; Schwartz et al. 2024; Schwarz et al. 2025). Bacteria-based liquid cultures have been described for a number of entomopathogenic nematodes, most of which utilized their associated symbiotic bacteria as the food source (Buecher and Popiel 1989; Lunau et al. 1993; Ehlers et al. 1998; Ehlers et al. 2000; Ferreira et al. 2014; Addis et al. 2016; Ferreira et al. 2016; Pérez-Campos et al. 2018; Dunn et al. 2019). Here, we describe a liquid culture method for *S. hermaphroditum* that we adapted and modified from the method used in *C. elegans*. We showed that the worms could thrive under these conditions and that these liquid cultures can produce infective juveniles (IJs), with or without their symbiont.

We modified the S Medium-based liquid culture method widely used in *C. elegans* research (Stiernagle 2006; Hibshman et al. 2021) by replacing *Escherichia coli* with either the non-symbiont bacteria *Comamonas aquatica* (Avery and Shtonda 2003; Watson et al. 2014), or the native symbiont *Xenorhabdus griffinae* (Tailliez et al. 2006). We showed previously that *S. hermaphroditum* could be cultured on lawns of the non-symbiotic bacteria *C. aquatica* (Cao et al. 2022; Rodak et al. 2024). When cultured on agar-based media, *C. aquatica* offers several advantages over the native symbiont *X. griffinae*, including a higher growth rate, better transparency of the worm, and an apparent lack of phase variation observed in *Xenorhabdus* species (Akhurst 1980). We therefore adapted *C. aquatica* as the primary food source used in this study, while also developing protocols for liquid culture of *S. hermaphroditum* using its native symbiont *X. griffinae* as the food source.

We cultured *S. hermaphroditum* on varying concentrations of *C. aquatica* or *X. griffinae* starting from fertilized embryos and recorded the lengths of the worms as they grew, the amount of time they took to reach adulthood and reproduce, and the time needed for the population to produce dispersal-stage IJ larvae. We found that at sufficiently high food concentrations (20% of concentrated *C. aquatica* suspension or 10% of concentrated *X. griffinae* suspension in S medium by volume. Concentrated bacteria suspension was obtained by resuspending overnight culture in S medium 1/10 of the original volume (10X concentrated). See methods for details.) *S. hermaphroditum* can be successfully cultivated to adulthood (Fig. 1A) and produce progeny (Fig. 1B and C). In multiple independent tests, we found that increasing the food concentration increased the growth rate of the worm (Fig. 1D and E) and decreased the amount of time needed to reproduce (Fig. 1B and C). For example, when starting with an average of 762 worms/mL (± 220 , $n = 6$), F₁ progeny were observed on day 6 when *C. aquatica* (concentrated) suspension was 20% of the culture medium (by volume), but on day 4 when the *C. aquatica* suspension was 50% or 100% of the culture medium (Fig. 1B). Similarly, in a separate experiment with *X. griffinae* as food, F₁ progeny were observed on day 5 when bacteria concentrations were at 10% but on day 4 at 20% (Fig. 1C). When food was resupplied daily (adding 25 μ L, 50 μ L, or 100 μ L of concentrated *C. aquatica* to the original culture), multigenerational reproduction was observed, with newly matured adults observed up until at least day 20, suggesting the feasibility of continuous culture.

Without subsequent food supplements, the worms eventually cleared the bacteria, and young larvae developed into infective juveniles (IJs) (Fig. 1F) at around day 8-9 (Fig. 1B and C). We observed that the IJs of *S. hermaphroditum* are ensheathed, with the J2 cuticle clearly observable under microscopy (Fig. 1F). Ensheatment of IJs is characteristic of many parasitic nematodes, including EPNs (Lee 2002), but this cuticle is readily lost in the free-living nematode model *C. elegans* (Cassada and Russell 1975). Ensheatment could explain our previous observation that the sensory neurons of *S. hermaphroditum* IJs are not surface-exposed, as assessed using DiI (1,1'-Diocetadecyl-3,3,3',3'-Tetramethylindocarbocyanine Perchlorate) staining (Garg et al. 2022).

In summary, we have developed a method for a liquid culture of *S. hermaphroditum* using either its native symbiont, the entomopathogenic bacterium *X. griffinae*, or, as a neutral option, the soil bacterium *C. aquatica*, as a food source. We believe this small volume liquid culture method could be a useful tool for the study of *S. hermaphroditum* biology in the laboratory and as a starting point for testing methods for large volume liquid culture.

Methods

Solid media nematode culture

The wild-type strain *Steinernema hermaphroditum* strain PS9179 (Cao et al. 2022), which was inbred from the isolate CS34 (Bhat et al. 2019), was cultured on Enriched Peptone Medium agar (EPM) on a lawn that had been seeded from an overnight culture of *Comamonas aquatica* as described in Rodak et al. (2024).

Liquid culture methods

S Medium (base medium)

The recipe of S Medium was based on Stiernagle (2006) and Hibshman et al. (2021) (referred to as S-complete). Briefly, for ~1 liter of the solution, add:

- (1) 1 liter of S-basal solution [per 1 L: 5.9 g of NaCl, 50 mL of 1 M KH_2PO_4 solution pH 6, add deionized water to 1 L and autoclave to sterilize. After autoclave, add 1 mL of Cholesterol solution (5 mg/mL in ethanol)]
- (2) 10 mL of Trace metal solution [per 1 L: 1.86 g of Disodium EDTA, 0.69 g of $\text{FeSO}_4 \cdot 7\text{H}_2\text{O}$, 0.2 g of $\text{MnCl}_2 \cdot 4\text{H}_2\text{O}$, 0.29 g of $\text{ZnSO}_4 \cdot 7\text{H}_2\text{O}$, 0.025 g of $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$, add deionized water to 1 L and autoclave to sterilize. After autoclaving, store in the dark at room temperature.]
- (3) 10 mL of 1M Potassium citrate pH 6 [per 1 L: 20 g of citric acid monohydrate, 293.5 g of Tri-potassium citrate monohydrate, add deionized water to 1 L and autoclave to sterilize.]
- (4) 3 mL 1M CaCl_2 solution
- (5) 3 mL 1M MgSO_4 solution

Concentrated bacteria (Food source)

Recently thawed *Comamonas aquatica* DA1877 (Avery and Shtonda 2003), and *Xenorhabdus griffinae* HGB2511 (Cao et al. 2022) were streaked onto Luria-Bertani (LB) plates, grown at 37°C and room temperature respectively, and stored either at 4°C (DA1877) or at room temperature in the dark (HGB2511). To prepare concentrated bacteria as the food source, LB medium (200 mL) in Erlenmeyer flasks (1 L size) was seeded with a single picked colony of bacteria and incubated overnight at 200 rpm at 37°C (DA1877) or at 30°C (HGB2511). The bacteria from the overnight culture were pelleted by centrifugation (7745 rcf, 20 minutes) at 4°C, the supernatant was removed, and the pellet was resuspended in S Medium at 10% of the original culture volume. The concentrated bacteria solution was then stored at 4°C until usage.

To prepare the bacteria-based liquid culture medium, the concentrated bacteria solution was warmed up to room temperature and diluted with S Medium to create the different concentrations of the medium used in this study. We do not expect either bacterium to have significant growth in S medium.

Bacterial-based liquid culture and analyses

In the liquid culture described in this study, worms were cultured in 24-well tissue culture plates incubated at 25°C on an orbital shaker platform moving at 60 rpm. Inside the incubator, the plates were placed in a plastic shoebox along with wet paper towels that were wetted regularly to maintain humidity. Each well contained 1 mL of culture medium.

To initiate the culture, gravid *S. hermaphroditum* adults grown on *Comamonas*/EPM plates were harvested to obtain embryos based on the method described in Rodak et al. (2024). Briefly, gravid adults were collected from the plates using M9 buffer into a 15 mL centrifuge tube. The worms were then pelleted by centrifugation to remove the supernatant. The worm pellet was then resuspended with water to obtain a total volume of 3.5 mL. 0.5 mL of 5 M NaOH and 1 mL of household bleach (8% available chlorine) were then added to the solution for a total volume of 5 mL. The solution was mixed by shaking and allowed to react for about 5 minutes, by which time most tissues other than the embryos should be visibly dissolved. In our experience, longer reactions substantially decreased the viability of the embryos and should preferably be avoided. The embryos were collected by centrifugation and washed 3 times, each with 10 mL of M9 buffer. After the final wash, the embryos were again pelleted and resuspended in a small volume of M9 before being transferred to liquid cultures. To estimate the number of P_0 worms, 100 μL of liquid cultures were transferred onto a solid medium DA1877/ EPM plates the day following the initiation of the culture, animals were counted and removed. The plates were checked again the next day for worms that were missed. Six wells were sampled, and the average was calculated. In continuous culture assay, 1 mL cultures containing 20% *C. aquatica* suspension were resupplied daily by adding 25 μL , 50 μL , or 100 μL of concentrated *C. aquatica* to the original culture.

In Fig. 1B and C, the presence of F_1 progeny was determined by the simultaneous presence of both gravid P_0 adults and newly hatched F_1 larvae. The identification of infective juveniles (IJs) was based on features common to the IJ or dauer stages of *Steinernema* and some other nematode species, including slim body shape, sealed mouth, and the presence of an unecdysed J2 cuticle. The observations were performed using a Zeiss Imager Z2 microscope, in which 10 μL of liquid *Steinernema* culture

was mounted on a 2% agarose pad, with 1 µL of 100 mM levamisole added to immobilize the animals. Images used in Fig. 1A and 1F were obtained through an Axiocam 506 mono and Zen 2 Blue software using the same configurations. In Fig. 1D and E, worms were sampled every 24 hours by transferring a small portion of the culture onto empty NGM (nematode growth media) (Brenner 1974) plates, where the images of worms were captured using an M2Bio hybrid stereo-compound fluorescence microscope. Worm length was measured using ImageJ (Fiji) (Schindelin et al. 2012) as described in Rodak et al. (2024).

Acknowledgements: We thank members of the Sternberg Lab for their support. We would especially like to thank Hillel Schwartz for his valuable suggestions and for the critical reading of the manuscript. The *C. aquatica* strain DA1877 was obtained from the Caenorhabditis Genetics Center (CGC), which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440).

References

- Addis T, Mijušković N, Strauch O, Ehlers RU. 2016. Life history traits, liquid culture production and storage temperatures of *Steinernema yirgalemense*. *Nematology* 18: 367-376. DOI: [10.1163/15685411-00002966](https://doi.org/10.1163/15685411-00002966)
- Akhurst RJ. 1980. Morphological and Functional Dimorphism in *Xenorhabdus* spp., Bacteria Symbiotically Associated with the Insect Pathogenic Nematodes *Neoaplectana* and *Heterorhabditis*. *Microbiology* 121: 303-309. DOI: [10.1099/0021287-121-2-303](https://doi.org/10.1099/0021287-121-2-303)
- Avery L, Shtonda BB. 2003. Food transport in the *C. elegans* pharynx. *J Exp Biol* 206(Pt 14): 2441-57. PubMed ID: [12796460](https://pubmed.ncbi.nlm.nih.gov/12796460/)
- Baugh LR, Kurhanewicz N, Sternberg PW. 2011. Sensitive and precise quantification of insulin-like mRNA expression in *Caenorhabditis elegans*. *PLoS One* 6(3): e18086. PubMed ID: [21445366](https://pubmed.ncbi.nlm.nih.gov/21445366/)
- Bhat AH, Chaubey AK, Shokoohi E, Mashela PW. 2019. Study of *Steinernema hermaphroditum* (Nematoda, Rhabditida), from the West Uttar Pradesh, India. *Acta Parasitologica*. 64: 720-737. DOI: [10.2478/s11686-019-00061-9](https://doi.org/10.2478/s11686-019-00061-9)
- Boyd WA, Cole RD, Anderson GL, Williams PL. 2003. The effects of metals and food availability on the behavior of *Caenorhabditis elegans*. *Environ Toxicol Chem*. 22: 3049-55. PubMed ID: [14713049](https://pubmed.ncbi.nlm.nih.gov/14713049/)
- Brenner S. 1974. The genetics of *Caenorhabditis elegans*. *Genetics*. 77: 71-94. PubMed ID: [4366476](https://pubmed.ncbi.nlm.nih.gov/4366476/)
- Buecher EJ, Popiel I. 1989. Liquid Culture of the Entomogenous Nematode *Steinernema feltiae* with Its Bacterial Symbiont. *J Nematol*. 21: 500-4. PubMed ID: [19287644](https://pubmed.ncbi.nlm.nih.gov/19287644/)
- Cao M. 2023. CRISPR-Cas9 genome editing in *Steinernema* entomopathogenic nematodes. *bioRxiv*: 10.1101/2023.11.24.568619. DOI: [10.1101/2023.11.24.568619](https://doi.org/10.1101/2023.11.24.568619)
- Cao M, Schwartz HT, Tan CH, Sternberg PW. 2022. The entomopathogenic nematode *Steinernema hermaphroditum* is a self-fertilizing hermaphrodite and a genetically tractable system for the study of parasitic and mutualistic symbiosis. *Genetics* 220: 10.1093/genetics/iyab170. PubMed ID: [34791196](https://pubmed.ncbi.nlm.nih.gov/34791196/)
- Cassada RC, Russell RL. 1975. The dauerlarva, a post-embryonic developmental variant of the nematode *Caenorhabditis elegans*. *Dev Biol*. 46: 326-42. PubMed ID: [1183723](https://pubmed.ncbi.nlm.nih.gov/1183723/)
- Corsi AK, Wightman B, Chalfie M. 2015. A Transparent Window into Biology: A Primer on *Caenorhabditis elegans*. *Genetics*. 200: 387-407. PubMed ID: [26088431](https://pubmed.ncbi.nlm.nih.gov/26088431/)
- Dunn MD, Belur PD, Malan AP. 2019. *In vitro* liquid culture and optimization of *Steinernema jeffreyense* using shake flasks. *BioControl*. 65: 223-233. DOI: [10.1007/s10526-019-09977-7](https://doi.org/10.1007/s10526-019-09977-7)
- Ehlers RU, Lunau S, Krasomil Osterfeld K, Osterfeld KH. 1998. Liquid culture of the entomopathogenic nematode-bacterium-complex *Heterorhabditis megidis*/*Photorhabdus luminescens*. *BioControl*. 43: 77-86. DOI: [10.1023/a:1009965922794](https://doi.org/10.1023/a:1009965922794)
- Ehlers RU, Niemann I, Hollmer S, Strauch O, Jende D, Shanmugasundaram M, et al., Burnell A. 2000. Mass Production Potential of the Bacto-Helminthic Biocontrol Complex *Heterorhabditis indica* - *Photorhabdus luminescens*. *Biocontrol Science and Technology*. 10: 607-616. DOI: [10.1080/095831500750016406](https://doi.org/10.1080/095831500750016406)
- Felix MA. 2006. *Osccheius tipulae*. *WormBook*: 1-8. PubMed ID: [18050438](https://pubmed.ncbi.nlm.nih.gov/18050438/)
- Ferreira T, Addison MF, Malan AP. 2014. *In vitro* Liquid Culture of a South African Isolate of *Heterorhabditis zealandica* for the Control of Insect Pests. *African Entomology*. 22: 80-92. DOI: [10.4001/003.022.0114](https://doi.org/10.4001/003.022.0114)
- Ferreira T, Addison MF, Malan AP. 2016. Development and population dynamics of *Steinernema yirgalemense* (Rhabditida: Steinernematidae) and growth characteristics of its associated *Xenorhabdus indica* symbiont in liquid culture. *J Helminthol*.

4/28/2025 - Open Access

90: 364-71. PubMed ID: [26156314](#)

Garg P, Tan CH, Sternberg PW. 2022. DiI staining of sensory neurons in the entomopathogenic nematode *Steinernema hermaphroditum*. *MicroPubl Biol*. 2022 PubMed ID: [35224464](#)

Hibshman JD, Webster AK, Baugh LR. 2021. Liquid-culture protocols for synchronous starvation, growth, dauer formation, and dietary restriction of *Caenorhabditis elegans*. *STAR Protoc*. 2: 100276. PubMed ID: [33490989](#)

Lam ABQ, Webster JM. 1971. Morphology and Biology of *Panagrolaimus-Tipulae N Sp* (Panagrolaimidae) and *Rhabditis-(Rhabditella)-Tipulae N Sp* (Rhabditidae), from Leatherjacket Larvae, *Tipula-Paludosa* (Diptera-Tipulidae). *Nematologica*. 17: 201-212.

Lee DL. 2002. *The biology of nematodes*. PubMed ID: [14134023](#)

Lunau S, Stoessel S, Schmidtpeisker AJ, Ehlers RU. 1993. Establishment of Monoxenic Inocula for Scaling-up *in-Vitro* Cultures of the Entomopathogenic Nematodes *Steinernema Spp* and *Heterorhabditis Spp*. *Nematologica*. 39: 385-399. DOI: [Doi 10.1163/187529293x00330](#)

Maupas E. 1900. Modes et formes de reproduction des nematodes. *Arch. Zool. Exp. Gen*. 8: 463-624.

Park HH, Jung Y, Lee SV. 2017. Survival assays using *Caenorhabditis elegans*. *Mol Cells*. 40: 90-99. PubMed ID: [28241407](#)

Perez Campos SJ, Rodriguez Hernandez AI, Del Rocio Lopez Cuellar M, Zepeda Bastida A, Chavarria Hernandez N. 2018. *In-vitro* liquid culture of the entomopathogenic nematode, *Steinernema colombiense*, in orbitally shaken flasks. *Biocontrol Science and Technology*. 28: 901-911. DOI: [10.1080/09583157.2018.1499872](#)

Rodak NY, Tan CH, Sternberg PW. 2024. An improved solid medium-based culturing method for *Steinernema hermaphroditum*. *MicroPubl Biol*. 2024. PubMed ID: [38344064](#)

Schindelin J, Arganda Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, et al., Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods*. 9: 676-82. PubMed ID: [22743772](#)

Schwartz HT, Tan CH, Peraza J, Raymundo KLT, Sternberg PW. 2023. Molecular identification of a peroxidase gene controlling body size in the entomopathogenic nematode *Steinernema hermaphroditum*. *Genetics* 226: 10.1093/genetics/iyad209. PubMed ID: [38078889](#)

Schwarz EM, Baniya A, Heppert JK, Schwartz HT, Tan CH, Antoshechkin I, et al., Dillman AR. 2025. Genomes of the entomopathogenic nematode *Steinernema hermaphroditum* and its associated bacteria. *bioRxiv*: 2025.01.09.632278. DOI: [10.1101/2025.01.09.632278](#)

Sommer RJ. 2006. *Pristionchus pacificus*. *WormBook*: 1-8. PubMed ID: [18050490](#)

Sommer RJ, Carta LK, Kim SY, Sternberg PW. 1996. Morphological, genetic and molecular description of *Pristionchus pacificus* sp n (Nematoda: Neodiplogastridae). *Fundamental and Applied Nematology*. 19: 511-521.

Stiernagle T. 2006. Maintenance of *C. elegans*. *WormBook*: 1-11. PubMed ID: [18050451](#)

Stock SP, Griffin CT, Chaerani R. 2004. Morphological and molecular characterisation of *Steinernema hermaphroditum* n. sp (Nematoda : Steinernematidae), an entomopathogenic nematode from Indonesia, and its phylogenetic relationships with other members of the genus. *Nematology*. 6: 401-412. DOI: [Doi 10.1163/1568541042360555](#)

Tailliez P, Pages S, Ginibre N, Boemare N. 2006. New insight into diversity in the genus *Xenorhabdus*, including the description of ten novel species. *Int J Syst Evol Microbiol*. 56: 2805-2818. PubMed ID: [17158981](#)

Watson E, Mac Neil LT, Ritter AD, Yilmaz LS, Rosebrock AP, Caudy AA, Walhout AJ. 2014. Interspecies systems biology uncovers metabolites affecting *C. elegans* gene expression and life history traits. *Cell*. 156: 759-70. PubMed ID: [24529378](#)

Funding: This research was supported by National Science Foundation (NSF) Enabling Discovery through Genomics (EDGE).

Supported by U.S. National Science Foundation (United States) 2128267 to PWS.

Author Contributions: Nathan Y. Rodak: investigation, formal analysis, methodology, visualization, writing - original draft. Chieh-Hsiang Tan: conceptualization, formal analysis, methodology, supervision, visualization, writing - original draft, writing - review editing. Paul W. Sternberg: conceptualization, funding acquisition, supervision, writing - review editing.

Reviewed By: Curtis Loer

WormBase Paper ID: WBPaper00068074

4/28/2025 - Open Access

History: **Received** February 18, 2025 **Revision Received** April 6, 2025 **Accepted** April 25, 2025 **Published Online** April 28, 2025 **Indexed** May 12, 2025

Copyright: © 2025 by the authors. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International (CC BY 4.0) License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Citation: Rodak NY, Tan CH, Sternberg PW. 2025. A small-scale bacterial-based liquid Culture Method for *Steinernema hermaphroditum*. microPublication Biology. [10.17912/micropub.biology.001549](https://doi.org/10.17912/micropub.biology.001549)